

SH80: What are the indications for debridement, antibiotics and implant retention (DAIR) in subacute and chronic shoulder PJI?

Liaison: Surena Namdari

Lead Delegate: Mark Falworth

Supportive Delegates: Fernando Santana; Gabor Skaliczki

Response: In the subacute/chronic PJI setting, DAIR may have a role in patients with a well-functioning well-fixed implant who are unable to undergo extensive revision surgery.

Strength of Recommendation: Limited

Delegate Vote: 49 (96%) agree; 0 disagree; 2 (4%) abstain

Rationale: A comprehensive literature search was performed in 2024 using databases: Medline, Embase, Web of Science, CINAHL, Scopus, Cochrane, Clinicaltrial.gov and PubMed to identify all studies on irrigation and debridement with implant retention (DAIR) when treating shoulder PJI. The search terms were Periprosthetic Joint Infection or Prosthesis-Related Infections, Shoulder or Shoulder Pain or Shoulder Joint or shoulder, Shoulder Joint or Arthroplasty, Replacement, Shoulder or Arthroplasty, Replacement or shoulder arthroplasty or Joint Prosthesis, Surgical Wound Infection or Shoulder Prosthesis or Anti-Bacterial Agents or intrawound shoulder or Vancomycin, Arthroplasty, Replacement, Shoulder/ or Prosthesis-Related Infections or Anti-Bacterial Agents or chronic shoulder periprosthetic joint infection or Shoulder Joint, debridement, antibiotics and implant retention, persistent positive culture or positive culture, Orthopaedic / orthopedic procedures or orthopaedic / orthopedic surgical protocols or Postoperative Complications. The systematic review software Rayyan was used to deduplicate the articles and for the literature screening process. *Inclusion criteria* for the search were English language articles, all papers include the shoulder arthroplasty, or Periprosthetic Joint Infection. *Exclusion criteria* were non-English language articles, animal studies, single case studies, case report studies, cancer, dentistry, knee, hip, ankle, spine, and elbow papers, however due to the low number of relevant publications found, relevant lower limb PJI publications were also reviewed.

Although debridement, antibiotics and implant retention (DAIR) for a well-fixed shoulder implant may have a role in the management of prosthetic joint infection (PJI), there are no prospective or randomised studies to clarify its role in the treatment of subacute or chronic shoulder PJI. Furthermore, an accepted definition of what constitutes a subacute or chronic infection is also lacking, making comparisons of available studies difficult.

The timing of treatment is perceived as being crucial for the successful management of PJI, as treatment recommendations will vary based on whether there has been enough time for the formation of the biofilm, following which component retention is discouraged [1,2]. The Infectious Diseases Society of America (IDSA) has therefore suggested that DAIR should only be considered within 30 days of the index surgery, or within 21 days of symptoms related to the infection, with any PJI noted beyond this period not being considered for DAIR [1].

However, the orthopaedic literature has used a range of timings when managing shoulder PJI, which causes some confusion with respect to DAIR treatment algorithms and recommendations. Using IDSA recommendations for what constitutes a subacute or chronic infection (onset < 30 days or < 21 days of symptoms), there are two studies that reveal very conflicting results, with one recording 80% failure (4 of 5 cases), whereas another demonstrated 100% eradication of infection (2 cases), but with one of the cases requiring three DAIR procedures [1,3,4]. Where considering only subacute infection, which has been

suggested as being between 3 and 12 months, 50% failure (3 of 6 cases) was noted in one study, and 100% failure (3 of 3 cases) in another [5,6,7]. In contrast, where chronic infection (> 12 months) was managed with DAIR, other studies have reported success rates as 50% (1 of 2 cases), 71% (12 of 17 cases) and 88% (7 of 8 cases) [5,8,9]. However, the variation in surgical technique for DAIR, including if there was any modular exchange, varied from study to study, such that direct comparison between the reports is problematic.

Therefore, the use of DAIR in the presence of a well-fixed implant with subacute or chronic PJI is unclear, however it may still have a role, particularly in a patient who cannot tolerate staged revision surgery.

References:

1. Osmon DR, Berbari EF, Berendt AR, Lew D, Zimmerli W, Steckelberg JM, Rao N, Hanssen A, Wilson WR. Diagnosis and management of prosthetic joint infection: clinical practice guidelines by the Infectious Diseases Society of America. Infectious Diseases Society of America. Clin Infect Dis. 2013 Jan;56(1):e1-e25.
2. Dennison T, Alentorn-Geli E, Assenmacher AT, Sperling JW, Sánchez-Sotelo J, Cofield RH. Management of acute or late hematogenous infection after shoulder arthroplasty with irrigation, débridement, and component retention. J Shoulder Elbow Surg. 2017 Jan;26(1):73-78.
3. Lemmens L, Geelen H, Depypere M, De Munter P, Verhaegen F, Zimmerli W, Nijs S, Debeer P, Metsemakers WJ. Management of periprosthetic infection after reverse shoulder arthroplasty. J Shoulder Elbow Surg. 2021 Nov;30(11):2514-2522.
4. Bdeir M, Dally FJ, Assaf E, Gravius S, Mohs E, Hetjens S, Darwich A. Periprosthetic Infections of the Shoulder Joint: Characteristics and 5-Year Outcome of a Single-Center Series of 19 Cases. Antibiotics (Basel). 2021 Sep 18;10(9):1125.
5. Sperling JW, Kozak TK, Hanssen AD, Cofield RH. Infection after shoulder arthroplasty. Clin Orthop Relat Res. 2001 Jan;(382):206-16.
6. Jacquot A, Sirveaux F, Roche O, Favard L, Clavert P, Molé D. Surgical management of the infected reversed shoulder arthroplasty: a French multicenter study of reoperation in 32 patients. J Shoulder Elbow Surg. 2015 Nov;24(11):1713-22.
7. Ortmaier R, Resch H, Hitzl W, Mayer M, Stundner O, Tauber M. Treatment strategies for infection after reverse shoulder arthroplasty. Eur J Orthop Surg Traumatol. 2014 Jul;24(5):723-31.
8. Kew ME, Mathew JJ, Wimberly AC, Fu MC, Taylor SA, Blaine TA, Carli AV, Dines JS, Dines DM, Gulotta LV. Outcomes after débridement, antibiotics, and implant retention for prosthetic joint infection in shoulder arthroplasty. J Shoulder Elbow Surg. 2024 Feb;33(2):e68-e78.
9. Coste JS, Reig S, Trojani C, Berg M, Walch G, Boileau P. The management of infection in arthroplasty of the shoulder. J Bone Joint Surg Br. 2004 Jan;86(1):65-9.

