

SH77. Should bone graft or cement be removed during treatment of acute shoulder PJI?

Mark Lazarus MD; Grant Garrigues MD; Robert Hudek MD

Methodology: A comprehensive literature review was performed to identify all studies on retained bone cement and/or bone graft during revision shoulder arthroplasty. Searches of PubMed and Google Scholar were performed through October 2024 for the MeSH major topic “arthroplasty” subheading “cement,” major topic “arthroplasty” and subheading “bone graft,” and combination of all field and MeSH terms “arthroplasty” “infection” “cement” “bone graft” and “revision.” Because of a paucity of studies specifically reporting shoulder arthroplasty available on this topic, the search was expanded to include studies for other joints. Inclusion criteria were all English studies (Level I-IV evidence) that reported on the influence of retained cement and/or bone graft during revision arthroplasty for prosthetic joint infection (PJI). When present in the report, specific outcomes of patients treated during the acute period (4 – 12 weeks) after PJI is reported. In the studies identified, it was impossible to differentiate those patients with acute PJI from hematogenous spread from those with subacute or chronic. Exclusion criteria were non-English language articles, nonhuman studies, retracted papers, case reports, review papers, studies with less than <10 patients in the sample size, studies without clinical follow-up/infection rates, and technique papers without patient data. PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) criteria were followed.

Answer: Unknown

Strength of Recommendation: Limited

Rationale:

There is one study that investigated the effect of cement retention during revision surgery for shoulder PJI on subsequent or persistent infection (1). This study did not differentiate acute vs. subacute or chronic patients. 16% of the 37 total patients had retained cement. Of the 10 patients with recurrent infection, only 1 had retained cement ($p=0.389$). The overall infection rate of those with retained cement was 7% as compared to 30% for those without retained cement. 48.6% of infections were *C. Acnes* and this group had a cure rate of 89%. Of the 4 infections with staph species (MRSA or MSSA), only 1 was cured. The distribution of offending bacterium in the retained vs. cleared-of-cement groups is not reported.

There are two other upper extremity studies, both in the elbow. One study reported on 57 elbows that underwent revision total elbow arthroplasty (TEA) for infection, with an infection eradication rate of 69%(2). The authors do define patients as acute (1), subacute (9), and chronic (42), but do not specifically report results by group. Overall, time from index surgery to first debridement did not statistically change recurrence rate ($p=0.3$). The risk of persistent infection was 3.3 times higher in those patients with retained cement ($p=0.04$) but the risk of continued infection was more related to infecting organism (Staph Epi, $p=0.06$). The second study analyzed 19 patients after revision TEA for infection (3). The time from index surgery to first revision was 7.3 years (range 0.1 – 24.7), but the exact distribution of patients operated on in the acute, subacute, or chronic periods was not reported. One patient was infected with *C. Acnes*. The overall infection eradication rate was 53%. Six patients (60%) with retained cement after first revision had persistent

infection as compared to 3 patients (33%) with full cement removal ($p=.370$). Three patients (30%) who had retained cement required allograft during final revision as compared to 6 patients (67%) who had complete cement removal ($p=.179$), suggesting that attempts at complete cement removal may be detrimental to subsequent bone integrity.

There are three studies in the hip arthroplasty literature that specifically address the influence of retained cement during revision for PJI on subsequent infection. One study analyzed 89 consecutive patients and reported a persistent infection rate of 7.9% after single stage revision with retention of cement (4). The procedure described is meticulous, including burring of any cement that was exposed to the joint fluid space. The authors do not define patients by acute, subacute, or chronic, but 42.7% were operated on within the first 6 months from index arthroplasty. Results based on time from index arthroplasty to revision were not reported. Only 2 patients were deemed infected with “propionobacterium” species, and both of these patients were in the cured cohort. Another study in the hip arthroplasty literature reviewed 15 patients with only 1 suffering persistent or recurrent infection but 2 patients having positive cultures at the time of subsequent surgery (5). Although months from index procedure to revision was reported (50, range 0.75 – 155), results were not differentiated based on acute vs. subacute/chronic presentation. The most common infecting organism was Staph A. and none of the patients were infected with cutibacterium. Contrary to these results, one final study in the hip literature reported on 10 patients who underwent revision with retained cement, with a successful eradication of infection in only 2 (6). Reaming or burring of the remaining cement mantle was not routinely performed. Only 1 patient was diagnosed and treated in the acute postoperative period, infecting organism Staph Epi, and this patient was one of the two cures.

There are no studies in any joint specifically investigating the effect of bone graft removal on eradication of infection after PJI.

References:

1. Schiffman CJ, Kane L, Khoo KJ, Hsu JE, Namdari S: “Does retained cement or hardware during 2-stage revision shoulder arthroplasty for infection increase the risk of recurrent infection?,” JSES (2024), epub ahead of print.
2. Martinez-Catalan N, Nguyen NT, Morrey ME, O’Driscoll SW, and Sanchez-Sotelo J: “Two-stage reimplantation for deep infection after total elbow arthroplasty,” *Shoulder & Elbow* 2022, Vol. 14(6) 668–676.
3. Schiffman CJ, Baker W, Kwak D, Ramsey ML, Namdari S, Austin LS: “High failure rate of 2-stage revision for the infected total elbow arthroplasty: a single institution’s experience,” *J Shoulder Elbow Surg* (2024) 33, S122–S129.
4. Fishley WG, Selvaratnam V, Whitehouse SL, Kassam M, Petheram TG: “Cement-in-cement revision of the femur in infected hip arthroplasty in 89 patients across two centres,” *Bone Joint J* 2022;104-B(2):212–220.
5. Morley JR, Blake SM, Hubble MJW, Timperley AJ, Gie GA, Howell JR: “Preservation of the original femoral cement mantle during the management of infected cemented total hip replacement by two-stage revision,” *J Bone Joint Surg Br* 2012;94-B:322–7.
6. Leijtens B, Sadeghi N, Schreurs BW, Rijnen WH: “Cement-within-cement revision of infected total hip replacement; disappointing results in 10 retrospective cases,” *Hip Int* 2016; 26 (1): 67-72.